

United States Department of Agriculture National Agricultural Statistics Service Michigan Field Office Cooperating with Michigan Department of Agriculture



MI-CW3008

Michigan Crop Weather

July 28, 2008

Varied Moisture

Six days were suitable for fieldwork during the week ending July 27, according to the USDA, NASS, Michigan Field Office. Precipitation varied from 0.02 inches in the west central Lower Peninsula to 0.79 inches in the east central Lower Peninsula. Average temperatures ranged from 3 degrees below normal in the southwestern and south central Lower Peninsula to 1 degree below normal in the eastern Upper Peninsula, and the northwestern and northeastern Lower Peninsula. Temperatures this past week were slightly below normal but still fairly warm throughout the state. Growers in many areas reported that there is a need for rainfall, while others are saying that there is too much moisture. A reporter in the central region stated, "We still continue to battle too much moisture in some areas, while others are getting dry. There is a huge difference in moisture levels within a 2-3 mile drive!' Harvest of wheat and hay is great in the drier areas, but difficult in others due to moisture levels. "Second cutting hay is hard to harvest as we have had foggy mornings and overcast afternoons. Some wheat is getting harvested in the area but they have had the same problem with moisture," a farmer in the southwest reported.

Field Crops

The varied moisture levels have presented a challenge for some farmers. Corn continued to grow well this past week and was tasseling in many fields. Most were optimistic about yield potential. Continue to monitor for pests such as cutworm and armyworm. Some of the soybeans that were having problems earlier were improved because of the sun and rains, but some fields still have yellow areas. At this stage, the majority of the crop is blooming and some are setting pods. A large amount of winter wheat was harvested this week and was approaching completion in many areas, but some were still harvesting as weather and moisture levels permit. Harvest of alfalfa is progressing. Rains and moisture slowed harvest in some areas while dry conditions were great for harvest in other areas. Dry beans stands were average and some reported damage due to excess moisture. The sugarbeet crop looked good. Growers were very optimistic about production potential. Oat harvest was just beginning in some areas, though many expected to begin this coming week. Barley remained in good condition.

Fruit

Apples grew to 2.8 inches and 47 mm in diameter in the southwest and northwest, respectively. In the southeast, where new strikes of fire blight infections were found, apples grew to 2.5 inches. In the Grand Rapids area, apples sized well and growers were encouraged to closely scout for insect pests. Harvest of midseason apricots was underway in the southwest and northwest. Blackberry harvest continued in the southwest. Cranberries were at pinhead and small green fruit in the southwest. Blueberry harvest was underway across the State. In the southeast, where berries have been slow to ripen on many farms, the demand for pick your own and ready pick berries was excellent. In the southeast, harvest of early variety peaches began; oriental fruit moths were prevalent. Pears were 35 mm in the northwest and 2 inches in the southwest. In the southeast, pears grew to 1.75 inches; fire blight and leaf spot infections were found. Plums were 27 mm in the northwest, and harvest of 'Early Golden' and 'Methley' began in the southwest. **Strawberry** re-growth from renovation was underway in the southwest; renovation was completed in the southeast. Sweet cherry harvest was underway in the west central; a large amount of abandonment due to excessive splitting was expected. In the northwest, harvest was in full swing, as fruit quality was a challenge with frost scars and cracking from varied rainfall. In the southwest, sweet cherry harvest was completed and post-harvest pruning began. Tart cherry harvest was underway in the west central and northwest; harvest neared completion in the southeast. Raspberry harvest began in the northwest and continued in the southwest and southeast. Grapes were at cluster tightening in the southwest and berry touch in the northwest.

Vegetables

Temperatures continued about normal but rainfall amounts were low, and irrigation was underway where it was available. The first flush of fern on full season asparagus fields were fully expanded and most growers put down their first fungicide cover. Cabbage harvest continued at a good pace. Foliar diseases continued to be a concern for carrots. Root diameter in fields for processing was one to one and a half inches. Celery planting was completed in all areas and harvest was on schedule. Cantaloup harvest began this week. Watermelons were continuing to size with harvest approaching. Powdery mildew began to appear in most pumpkin fields. Early cucumber, yellow squash, and zucchini fields were finished and were being removed with some being replanted. Early virus symptoms were showing up. Onions continued to progress and bulbs were sizing at one to one and a half inches in diameter. Growers report normal insect and disease pressure with crop development possibly a little behind normal. Peppers and eggplants continued to grow well with few problems. Potatoes were ready earlier than expected and peas were almost done. Butternut and winter squash for processing were growing well with first bloom appearing. Summer snap bean plantings continued to grow normally in most areas. Sweet corn harvest just began and volume has increased as more fields come into production. There were no major problems reported. Harvest of market tomatoes were in full swing with good quality. Initial harvest of processing tomatoes is approaching. Pickle harvest presented a challenge as wet and warm weather has pushed pickle size and slowed harvesting operations.

Soil moisture for week ending 07/27/08

Stratum	Very short	Short	Adequate	Surplus		
	Percent	Percent	Percent	Percent		
Topsoil	9	19	67	5		
Subsoil	6	20	70	4		

Crop condition for week ending 07/27/08

Crop condition for week ending 07/27/08								
Crop	Very poor	Poor	Fair	Good	Excellent			
	Percent	Percent	Percent	Percent	Percent			
All Hay	2	4	22	54	18			
Barley	1	6	29	62	2			
Corn	2	4	16	46	32			
Dry beans	8	13	30	36	13			
Oats	1	4	22	55	18			
Pasture	1	6	32	47	14			
Soybeans	3	10	27	46	14			
Winter Wheat	1	4	24	46	25			

Crop progress for week ending 07/27/08

Crop	This week	Last week	Last year	5-year average	
	Inches	Inches	Inches	Inches	
Corn, height	70	70 56 6		65	
	Percent	Percent	Percent	Percent	
All hay, second cutting	63	50	71	64	
All hay, third cutting	6	NA	9	6	
Blueberries, harvested	36	27	33	33	
Corn, silked	68	28	70	61	
Corn, dough	2	NA	4	2	
Dry beans, blooming	46	10	43	46	
Dry beans, setting pods	5	NA	NA	NA	
Oats, turning yellow	78	58	82	75	
Oats, harvested	6	2	21	14	
Peaches, harvested	15	4	18	10	
Potatoes, harvested	1	NA	NA	NA	
Soybeans, blooming	79	64	79	70	
Soybeans, setting pods	36	6	40	30	
Tart cherries, harvested	60	26	67	73	
Winter wheat, harvested	85	47	95	77	

Michigan Weather Summary for Week Ending 07/27/08 ¹

		Temperature		Cumulative growing degree days ²		Precipitation						
Station	Maximum	Minimum	Departure from normal	2008	2007	Normal	This week	Last two weeks	Last four weeks	Since April 1	Norn Since	For
			поппа						WEEKS		April 1	month
Ironwood	81	43		1,040	1,376		0.33	1.49	2.60	11.16		
Marquette	80	48		973	1,299		0.26	1.60	2.71	13.19		
Stephenson	84	47		1,280	1,435		0.02	1.58	3.38	13.03	12.27	2 12
Western UP	84	39	-2	1,070	1,348	1,117	0.36	1.44	2.50	11.50	12.27	3.42
Cornell	86	48		1,123	1,364		0.38	0.98	2.61	11.89		
Sault St Marie	79	54		976	1,235		1.04	1.60	3.34	11.93		
Eastern UP	86	45	-1	983	1,250	941	0.53	1.18	2.52	12.28	11.42	3.19
Beulah	82	52		1,363	1,485		0.43	0.78	1.74	15.60		
Lake City	82	51		1,309	1,436		0.20	0.87	1.53	17.03		
Old Mission	82	53		1,290	1,404		0.23	0.77	1.32	10.14		
Pellston	80	52		1,251	1,376		1.07	2.02	3.01	12.80		
Northwest	82	48	-1	1,261	1,389	1,271	0.37	1.11	1.80	13.16	10.99	2.99
Alpena	83	54		1,300	1,405		0.05	3.57	4.36	13.03		
Houghton Lake	81	51		1,358	1,436		0.46	1.58	1.95	14.73		
Rogers City	81	53		1,281	1,448		0.37	2.41	2.92	14.10		
Northeast	83	51	-1	1,318	1,420	1,226	0.38	2.62	3.17	13.90	10.99	3.18
Fremont	84	51		1,449	1,596		0.00	0.15	2.52	11.92		
Hart	82	50		1,388	1,507		0.00	0.19	3.68	14.03		
Muskegon	83	58		1,427	1,607		0.01	0.28	3.14	11.69		
West Central	84	50	-2	1,402	1,543	1,398	0.02	1.02	3.72	14.60	10.97	2.54
Alma	84	51		1,535	1,683		0.28	0.69	1.18	8.55		
Big Rapids	84	50		1,483	1,489		0.37	0.71	1.74	11.05		
Central	84	50	-2	1,493	1,584	1,463	0.41	0.83	1.87	9.40	11.83	2.97
Bad Axe	82	58		1,495	1,520		0.88	2.11	4.14	13.29		
Pigeon	83	56		1,444	1,484		0.40	1.34	2.48	9.55		
Saginaw	83	54		1,596	1,679		0.65	1.60	2.74	11.56		
Standish	84	53		1,404	1,486		1.50	2.78	3.66	12.05		
East Central	84	53	-1	1,460	1,546	1,448	0.79	1.98	3.16	11.25	10.84	2.92
Fennville	83	55		1,455	1,598		0.11	0.86	2.10	16.58		
Grand Rapids	84	58		1,684	1,812		0.01	0.25	4.40	15.44		
Holland	84	55		1,670	1,769		0.00	0.70	4.08	17.26		
South Bend, IN	84	57		1,747	1,913		0.06	0.87	2.92	10.92		
Watervliet	83 86	55 49	-3	1,594	1,745 1,736	1,576	0.22 0.19	1.36	3.97 4.29	13.92 15.47	12.68	3.09
Southwest	86	49	-3	1,613	1,/36	1,3/6	0.19	1.16	4.29	13.47	12.08	3.09
Belding	NA	NA		1,491	1,605		0.00	0.07	2.17	9.61		
Coldwater	82	51		1,611	1,636		0.00	0.61	3.74	11.22		
South Central	83 NA	53 NA	-3	1,639 1,587	1,697 1,678	1,564	0.05 0.21	0.95 1.00	3.07 3.77	11.38 11.90	12.45	3.22
	0.5	50		1.700	1.020		0.22	0.00	2.20	10.21		
Detroit Flint	85 84	58 53		1,798 1,689	1,830 1,668		0.22 0.55	0.80 1.86	3.39 3.29	10.31 10.58		
Romeo	84	58		1,570	1,603		0.00	2.09	3.29	10.58		
Tipton	85	56		1,627	1,653		0.00	0.36	6.22	15.87		
Toledo, OH	88	56		1,775	1,812		0.00	0.93	5.39	14.63		
Southeast	88	52	-2	1,639	1,696	1,539	0.34	1.15	4.40	13.08	12.13	3.11

Issued by the USDA, NASS, Michigan Field Office in cooperation with the U.S. Department of Commerce, Michigan State University's Cooperative Extension Service, Agricultural Meteorologist, Department of Geography, and Crop Advisory Team ALERTS.

Growing degree days (GDD) is the sum of daily mean temperatures minus 50 per day, 86 maximum and 50 minimum. The GDD is accumulative from April 1.